

# Meeting the Mental Health Needs of Children

*Jane E. Kavanagh, Elizabeth Brooks, Susan Dougherty, Marsha Gerdes, James Guevara, and David Rubin*

## EXECUTIVE SUMMARY

Recent estimates suggest that 10 to 20 percent of children have a diagnosable mental health disorder<sup>1, 2</sup> and 40 to 80 percent of children with mental health problems do not receive the services they need.<sup>1, 3, 4</sup> Despite robust evidence indicating the benefits of timely prevention, detection, and intervention, physical and mental health systems continue to miss early opportunities to improve outcomes for these children.

Recent health reform legislation including the Mental Health Parity and Addiction Equity Act of 2008 and the Patient Protection and Affordable Care Act of 2010 (PPACA, as modified by the Health Care and Education Reconciliation Act of 2010) have made important strides

in establishing mental health needs as equivalent with physical health needs. With the full implementation of these laws, all qualified health plans will be required to provide mental health services, including behavioral health treatment, at parity with medical benefits.<sup>5-7</sup> In order to optimize the impact of these provisions on children's mental health, federal agencies, states, and private payers must establish care and reimbursement standards that promote a children's mental health environment more reflective of the evidence on best practices.

Through an examination of the evidence surrounding key issues in children's mental health, this brief proposes policy actions to improve outcomes for children and their families.

## EVIDENCE

Providing prevention-focused intervention for children who exhibit behaviors that are predictive of mental health disorders can improve outcomes.

Many children's mental health systems rely on unproven interventions.

Structural barriers limit the availability and accessibility of children's mental health services.

## ACTION

Prevention-focused interventions in children's mental health should be included in the essential benefits package developed under PPACA.

States should make use of legislative and regulatory measures to require public and private insurance reimbursement for **prevention-focused** interventions in children's mental health.

States should make use of legislative and regulatory measures to require public and private payers to reimburse for **evidence-based** interventions in children's mental health.

States should take a more active role in managing the implementation and evaluation of children's mental health services.

States should make use of legislative and regulatory measures to support the **integration** of mental health and primary care services.

State, city, and county health and mental health systems should promote the integration of mental health and primary care services by developing the flexibility to blend and braid funding streams.<sup>1</sup>

Federal and state programs should fund the demonstration and evaluation of innovative mental health delivery models.

## EVIDENCE TO ACTION FINDINGS

**1 EVIDENCE:** Providing prevention-focused intervention for children who exhibit behaviors that are predictive of mental health disorders can improve outcomes.

Most children's mental health problems are left untreated until they reach the severity of a diagnosable disorder.<sup>8</sup> However, children usually exhibit behaviors that are predictive of the development of a disorder well before they receive a diagnosis.<sup>2</sup> Children exhibiting such predictive behaviors are up to five times more likely to develop full mental health disorders than their non-symptomatic peers.<sup>9-11</sup> If these children receive appropriate intervention at an early stage, it can prevent the onset of a diagnosable mental health disorder and improve emotional, behavioral, and cognitive outcomes.<sup>2, 2, 12-18</sup> Additionally, in several studies, the academic, behavioral, and health benefits of prevention-focused interventions have lasted more than 10 years after the intervention.<sup>17</sup> This evidence indicates the need for a shift towards prevention-focused intervention in children's mental health.

The current limited availability of prevention-focused intervention stems partly from the financing structure of children's mental health services. In most states, providers are required to submit diagnoses from the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* to be reimbursed for services provided. This results in some providers (up to 70 percent) using alternate diagnoses and often inflating the severity of a child's problems in order to provide or refer to services. When considering using alternate diagnoses, 65 percent of providers surveyed report doing so to obtain services for a patient, and 73 percent report doing so particularly for children with predictive behaviors that do not reach diagnostic severity.<sup>19</sup> This practice is not without potential adverse effects, including the potential stigmatization that results from a child being labeled with a mental health disorder.<sup>20</sup> Furthermore, such labels often serve as pretext for the prescription of psychotropic medications. This includes the use of atypical antipsychotics, which have been prescribed off-label in children increasingly in recent

years despite emerging evidence of adverse effects.<sup>21</sup> Alternately, a provider, knowing the unlikelihood of reimbursement or the dangers of stigmatization, may delay intervention for a child whose symptoms are not severe enough to warrant a diagnosis.<sup>19</sup> This practice is equally problematic, as children with predictive behaviors are more likely to develop a diagnosable mental health disorder and because timely intervention is very effective among this group.<sup>9-13, 15, 17, 22</sup> In either of these cases, the child with predictive behaviors is unlikely to receive appropriate prevention-focused intervention.

**ACTION:** Prevention-focused interventions in children's mental health should be included in the essential benefits package developed under PPACA.

Starting in 2014, PPACA will require all qualified health plans<sup>23</sup> and individual and small group plans<sup>24</sup> to provide a package of essential benefits including mental and behavioral health services.<sup>25</sup> Additionally, PPACA requires that all qualified plans<sup>26</sup> and individual plans<sup>27</sup> provide these services at parity with medical benefits. The ongoing process of defining the benefits that will be included in these packages provides great opportunity to reshape the children's mental health system to better reflect the known benefit of timely prevention-focused intervention. In defining the mental health services included in the essential benefits packages, prevention-focused interventions should be prioritized.

**ACTION:** States should make use of legislative and regulatory measures to require public and private insurance reimbursement for **prevention-focused** interventions in children's mental health.

In the interim—before the full implementation of PPACA—states should mandate reimbursement for prevention-focused interventions to support availability of and access to such services for children. Although Medicaid policy does not generally support intervention for children without a diagnosis, many states have worked within the framework of the current system

<sup>1</sup> In both blending and braiding of funds, the provision of services is supported by multiple systems; however, blending funds combines funds into a single pool while braided funds remain separately linked to the supplying administrative system.

<sup>2</sup> Effect sizes of 0.24 to 0.93

to expand services to children exhibiting predictive behaviors.<sup>28</sup> As of 2000, North Carolina Medicaid allows six visits annually for mental health services without a diagnosis for children under 21 years of age. These visits can include evaluation and individual or group therapy sessions.<sup>29</sup> Similarly, Washington state passed legislation in 2007 providing up to 20 mental health visits through a fee-for-service network for children who do not meet criteria for a diagnosable disorder.<sup>28</sup> While increased access to these types of clinic treatment is an important step in prevention, states should require public and private insurers to empower mental health professionals to extend a variety of clinic and non-clinic preventive services to children who meet a well-defined set of risk factors.

Though such change could increase the upfront burden on the children's mental health system, unnecessary treatment might be prevented by ensuring that fewer children receive more serious mental health diagnoses. As long as regulations are in place to ensure that treatment is given only when appropriately indicated, this could lead to cost savings over time.<sup>1, 30</sup> This could also result in children receiving more appropriate treatment, and, most importantly for both child outcomes and cost-effectiveness, lead to fewer children with diagnosable mental health disorders as they age.

Given the paucity of evidence in this area, efforts to implement prevention-focused interventions should be tracked and evaluated to determine with certainty whether these modifications result in better outcomes for children and are more cost-effective over time.

**2 EVIDENCE:** Many children's mental health systems rely on unproven interventions.

Ensuring best outcomes for children with mental health problems requires not only timely intervention but also the use of interventions that have proven effective. Despite the existence of evidence-based interventions for the treatment and prevention of a variety of children's mental health problems, many states, cities, and counties continue to use unevaluated interventions and even use interventions that have been proven ineffective.<sup>17, 31-34</sup> The most promising behavioral health interventions focus

on addressing behaviors exhibited by children as well as caregiver responses to these behaviors. A review of 130 prevention-focused studies demonstrated that children receiving behavioral or cognitive-behavioral interventions had better outcomes than 70 percent of those receiving other types of interventions or no interventions.<sup>13</sup> Furthermore, several studies have found that usual care approaches to children's mental health have little to no benefit to children.<sup>17, 35, 36</sup>

The limited dissemination and implementation of many of the most promising evidence-based interventions is partly due to current reimbursement practices that inadequately fund core characteristics of these programs. For instance, although children's mental health research strongly indicates the benefit of treating both caregiver and child as "patient,"<sup>15, 28, 32, 37</sup> private and public insurance plans often do not reimburse for such "family-focused" treatment. Defining the child as the sole patient, "individual therapy" is usually reimbursed, while "family therapy" is covered less often. This issue is further complicated when seeking reimbursement for caregiver-only sessions as part of a child's treatment.<sup>38</sup> Additionally, many insurance plans limit the number of mental health sessions that are reimbursable in any given year, which is sometimes fewer than the number indicated for evidence-based programs.<sup>39-41</sup> Providers are therefore forced to seek repeated reauthorization in order to follow treatment protocols that have proven most beneficial for child outcomes. These strict reimbursement requirements may interrupt the treatment course or even discourage some providers from offering or referring to evidence-based programs despite their proven efficacy.

**ACTION:** States should make use of legislative and regulatory measures to require public and private payers to reimburse for **evidence-based** interventions in children's mental health.

To support the use of evidence-based interventions, states should require public and private insurance plans to reimburse for key characteristics of these interventions. Primarily, insurers must expand their definition of "patient" to cover caregivers of children when appropriate. "Family therapy" and "multifamily group therapy" should be reimbursed comparable to "individual therapy." Insurance plans should also reimburse for services to

caregivers without the child present when the services are geared towards improving child outcomes. Some states have already taken the lead on this issue. Hawaii, Vermont, California, Colorado, and Arizona provide models for the billing of evidence-based ‘family-focused’ interventions.<sup>42</sup> Additionally, provisions must be built into public and private insurance plans to ensure that reimbursement for evidence-based interventions can be authorized for the duration of treatment. Removing these reimbursement-related barriers is a crucial step in increasing the dissemination and implementation of

evidence-based interventions. [See Table 1 for examples of evidence-based interventions]

**ACTION:** States should take a more active role in managing the implementation and evaluation of children’s mental health services.

To ensure the quality of care children receive, states should be more active in managing the implementation and evaluation of children’s mental health services. Coordinated management at the state level could

**TABLE 1. SELECTED EVIDENCE-BASED INTERVENTIONS IN CHILDREN’S MENTAL HEALTH**

Intervention	Treatment Focus & Aims	Outcomes
<b>Parent-Child Interaction Therapy (PCIT)</b> <a href="http://www.pcit.org">www.pcit.org</a>	<b>Age:</b> 2-7 years and their caregivers; recent adaptations for ages 8-12 <b>Treatment Focus:</b> Externalizing behavior problems	<b>Reduced:</b> • Re-reports of physical abuse <b>Improved:</b> • Parenting skills and attitudes • Child behaviors
<b>Trauma-Focused Cognitive Behavioral Therapy (TF-CBT)</b> <a href="http://tfcbt.musc.edu">tfcbt.musc.edu</a>	<b>Age:</b> 3-18 years and their caregivers <b>Treatment Focus:</b> Emotional/behavioral problems resulting from child sexual abuse; adaptations for use with other traumatic experiences	<b>Reduced:</b> • PTSD symptoms • Self-reported fear and anxiety • Symptoms of depression <b>Improved:</b> • General functioning • Positive parenting skills • Parent and child coping skills
<b>CBT for Depression: Coping with Depression for Adolescents (CWDA)</b> <a href="http://www.kpchr.org/public/acwd/acwd.html">www.kpchr.org/public/acwd/acwd.html</a>	<b>Age:</b> 12-18 years and their caregivers <b>Treatment Focus:</b> Depression and/or dysthymia	<b>Reduced:</b> • Symptoms of depression • Development of diagnosable depressive disorders
<b>CBT for Anxiety: Coping Cat</b>	<b>Age:</b> 6-17 years and their caregivers <b>Treatment Focus:</b> Anxiety	<b>Reduced:</b> • Symptoms of anxiety • Symptoms of associated depression <b>Improved:</b> • Coping skills
<b>The Incredible Years (IY)</b> <a href="http://www.incredibleyears.com">www.incredibleyears.com</a>	<b>Age:</b> Birth-12 years and their caregivers or teachers <b>Treatment Focus:</b> Externalizing behavior problems	<b>Reduced:</b> • Parental depression • Child behavior problems <b>Improved:</b> • Parental positive affect • Effective parenting techniques • Child social and emotional competence

\*This table highlights some promising evidence-based interventions in the field of child and adolescent mental health. It is not an exhaustive list of all interventions with a strong evidence base. The following resources offer more information:

- The California Evidence-Based Clearinghouse for Child Welfare: [www.cebc4cw.org](http://www.cebc4cw.org)
- Blueprints for Violence Prevention: [www.colorado.edu/cspv/infhouse/index.html](http://www.colorado.edu/cspv/infhouse/index.html)
- SAMHSA’s National Registry of Evidence-Based Programs and Practices: [www.nrepp.samhsa.gov](http://www.nrepp.samhsa.gov)
- The Office of Juvenile Justice and Delinquency Prevention’s Model Programs Guide: [www2.dsgonline.com/mpg](http://www2.dsgonline.com/mpg)

strengthen the evidence base surrounding new and existing interventions and ensure continued efficacy once interventions are implemented. This charge is not simply for the identification of evidence-based practice, but a larger commitment to understanding how to adapt interventions for the populations they serve. Such an approach appreciates the full ecology (e.g. political, social, familial) that influences the success or failure of a mental health intervention, not just the intervention and outcomes in isolation.<sup>43</sup>

The complexities of state mental health systems resist a one-size-fits-all approach to reform. However, states that have undertaken efforts to implement evidence-based interventions have developed various strategies involving regulation, training, evaluation, and financing that could serve as models for other states. For instance, New York has established year-long training on evidence-based interventions, altered the clinic rate structure to incentivize the use of evidence-based interventions, and incorporated federal dollars into program evaluation. Hawaii has implemented statewide training programs, convened best practice conferences, and published practice guidelines. Oregon took a regulatory approach and passed legislation requiring 75 percent of its public mental health services to be evidence-based by 2008, while Michigan developed systems for outcome data and evaluation to allow the data to drive the reform.<sup>44, 45</sup> These state-level reform efforts should be evaluated to inform best practices in future large scale evidence-based implementation efforts.

While each state must tailor its approach to its particular needs and the structure of its children's mental health system, states should consider the following approaches for using regulation, training, evaluation, and financing to move towards evidence-based intervention in children's mental health:

1. Prioritize and standardize evidence-based programs for funding except where programs are in demonstration as a pilot phase.
2. Provide funds for ongoing personnel training and evaluation to increase capacity and ensure that interventions are delivered with fidelity to protocol, whether in the public or private sector. This could involve embedding training in evidence-based interventions into licensing requirements for mental

health professionals or requiring professionals to be certified in specific interventions in order to receive reimbursement for services rendered.

3. Seek federal demonstration dollars for development and evaluation in areas where interventions are lacking and to ensure the continued efficacy of programs once widely implemented. (Notably, the recent health reform legislation establishes a federal grant program designed specifically to fund the delivery of evidence-based prevention services.)

### 3 EVIDENCE: Structural barriers limit the availability and accessibility of children's mental health services.

A successful shift to a prevention-focused and evidence-based approach in children's mental health is also contingent on addressing barriers to availability and access. The dissemination and implementation of evidence-based interventions will improve outcomes only if the programs reach the children who need them. As mental health needs have risen over recent decades, there has been no parallel increase in services.<sup>18</sup> Shortages in resources, including a lack of professionals trained in children's mental health, have left many communities without access to quality mental health services.<sup>15, 46-49</sup>

While the data surrounding shortfalls in mental health services are limited, reports consistently identify a disparity between need and availability. Between 40 to 80 percent of children are reported to not receive the services they need.<sup>1, 3, 4</sup> Even after being accepted for a mental health evaluation, 30 to 60 percent never attend an appointment.<sup>2, 50, 51</sup> Additionally, 29.3 percent of caregivers who reported their child had a need for special therapeutic or educational services, equipment, or counseling had difficulty in accessing the needed services.<sup>52</sup> Notably, rates of unmet mental health needs are highest among Latino, African-American, economically disadvantaged, and uninsured children<sup>3, 53</sup> and up to 75 percent of children in the child welfare system have been reported not to receive the mental health services they need.<sup>54</sup>

The factors contributing to underutilization of services are complex. Families most commonly attribute the limited use of mental health services to a lack of knowledge



about available services, lack of availability, lack of transportation, lack of understanding about the purpose and mechanism of treatment, financial constraints, children's mental health professional shortages, stigma related to mental health disorders, and concerns about the use of psychotropic medications.<sup>48, 55, 56</sup> Additionally, 40 to 60 percent of families that begin treatment terminate it prematurely, with many of them reporting similar barriers as a reason for termination.<sup>1, 57</sup>

Creative solutions are needed to increase the availability of children's mental health services as well as to address access barriers for children and families. While there are many possible approaches to these issues, including increased training in mental health for medical and non-medical personnel working with children and families or co-locating mental health services in non-traditional settings such as community centers, more evaluation and evidence is needed to establish the potential impact of these approaches.

One innovation targeting these issues with promising results is the integration of mental health and primary care services. Early studies with adults have shown integrated services to treat mental health problems more effectively than standard care.<sup>58</sup> Integration of mental health and primary care services for children has been reported to be especially effective at "capturing" children with mental health problems because these children are more frequent users of primary care and are more likely to have numerous medical conditions than children without mental health problems.<sup>59</sup> In one study, when referred to a mental health provider located outside of a primary care office, only 60 percent of families sought these services within six months, while more than 80 percent of these families had returned to the primary care pediatrician's office for medical reasons during this time.<sup>60</sup> Providers also report increased provider satisfaction and decreased stigmatization and access barriers for families when mental health and primary care services are integrated.<sup>61</sup>

Some existing models for the integration of services include co-locating mental health professionals in existing pediatric primary care settings and establishing telemedicine mental health consultation. Massachusetts has used telephone consultations to expand access to approximately 95 percent of children in the state. After an initial phone consultation, the child receives appropriate intervention or connection to community resources.<sup>62</sup>

When mental health and primary care services are integrated in these ways, providers report greater frequency of consultations and referrals for mental health services and increased satisfaction with the process.<sup>62, 63</sup>

Despite the potential benefit of integrating mental health and primary care services for children, current reimbursement practices largely do not support this approach.<sup>18, 64</sup> Under many insurance plans, the provision of multiple services by different practitioners within the same provider organization on the same day is not allowed.<sup>65</sup> Outpatient consultation to a primary care provider is also often not reimbursed. Similarly, there is often minimal or no reimbursement for services that are not face-to-face with the patient (the child), even though billing codes exist to categorize many of these services.<sup>31, 66</sup> Accordingly, efforts to coordinate and integrate services such as appointments between clinicians and caregivers to discuss a child's mental health, discussions of care with teachers and schools, and conferencing between interdisciplinary mental health teams are often not financially supported.<sup>65, 66</sup>

**ACTION:** States should make use of legislative and regulatory measures to support the **integration** of mental health and primary care services.

States should require public and private insurers to support the integration of children's mental health and primary care services. Primarily, states must remove restrictions on services provided to a child on a single day at the same provider organization. These rules prevent a primary care pediatrician and a mental health professional working in the same provider organization from coordinating the care of a child in a timely and convenient way. This is an unnecessary barrier for families trying to access services for their child. States should also expand reimbursement for activities that coordinate care such as care plan oversight meetings and teleconferencing between providers. Removing these restrictions is an essential step to promoting integrated, prevention-focused intervention in children's mental health and primary care. Notably, telehealth services for the treatment of behavioral health problems was highlighted as an example of the types of programs to be tested by the newly created Center for Medicare and Medicaid Innovation.<sup>67</sup>

**ACTION:** State, city, and county health and mental health systems should promote the integration of mental health and primary care services by developing the flexibility to blend and braid funding streams.

Recognizing the limited financial resources available for mental health systems, states should be more proactive in blending and braiding funding streams across systems to support mental health efforts. In both blending and braiding of funds, the provision of services is supported by multiple systems; however, blending funds combines funds into a single pool while braided funds remain separately linked to the supplying administrative system. Blending allows for greater flexibility, as the funds can be distributed from a single source to cover a variety of children’s mental health expenses while braiding requires more maneuvering of funds from different streams to cover approved expenses but allows for greater tracking and accountability of monies.<sup>68</sup> Reflecting the far-reaching benefits of timely and effective interventions in children’s mental health, funds from children’s mental health, education, child welfare, delinquency and crime prevention, and substance abuse streams should be combined in these ways to maximize the resources available to provide children’s mental health services. States where these practices are already in place include Nebraska, Ohio, Wisconsin, Vermont, Michigan, and Indiana.<sup>42</sup> A collaborative funding effort will also promote a multi-system, integrated approach to meeting the needs of children with mental health problems.

**ACTION:** Federal and state programs should fund the demonstration and evaluation of innovative mental health delivery models.

Currently, while it would seem that integrated models of service delivery, whether in primary care or other locations, could alleviate challenges around availability of and access to mental health services, such models are still in their infancy and are not widely disseminated, implemented, or evaluated. We need to better understand whether such models can significantly reduce disparities in access to and utilization of mental health services in underserved communities. Therefore, state and federal agencies should fund the demonstration and evaluation of innovative delivery models to improve availability and accessibility of mental health services while retaining a commitment to evidence-based approaches in children’s mental health.

**CONCLUSION**

Children’s mental health systems face the challenge of serving the diverse mental health needs of a growing number of children while capacity and funding do not keep pace. This brief presents evidence-driven actions to improve outcomes for children by moving towards a mental health system focused on prevention, evidence-based intervention, integrated mental health and primary care services, and ongoing evaluation.

**PolicyLab’s Work On Children’s Mental Health**

<p><b>Enhanced Parenting For Depressed Caregivers</b> The intervention aims to reduce symptoms of depression in caregivers of young children through a primary care-based parental screening and intervention using the Incredible Years model.</p>
<p><b>CSAW: Pilot Intervention</b> The intervention works with families in the child welfare system to reduce behavioral problems, decrease placement moves, and mitigate caregiver stress. Two evidence-based interventions, Parent-Child Interaction Therapy and Child Adult Relationship Enhancement training, are co-located in foster care agencies through partnership with the City of Philadelphia’s Department of Human Services and Department of Behavioral Health.</p>
<p><b>Translating Evidence-Based Developmental Screening</b> This project aims to improve the appropriate identification and treatment of early developmental and behavioral problems in children by examining the feasibility, adaptability, and effectiveness of standardized developmental screening in pediatric practices.</p>
<p><b>Evaluating the Nurse Family Partnership Program</b> The Nurse Family Partnership is a prenatal home visitation program that has been found to positively impact a variety of maternal and child outcomes. PolicyLab’s evaluation of the program in Pennsylvania aims to assess the impact on five specific outcome measures to inform practice improvement initiatives and dissemination efforts.</p>

## REFERENCES

1. U.S. Department of Health and Human Services. *Mental Health: A Report of the Surgeon General*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health;1999.
2. Department of Health and Human Services. *U.S. Public Health Service, Report of the Surgeon General's Conference on Children's Mental Health: Developing a National Action Agenda*. Washington, DC;2000.
3. Kataoka SH, Zhang L, Wells KB. Unmet need for mental health care among U.S. children: Variation by ethnicity and insurance status. *Am J Psychiatry*. 2002;159:1548-1555.
4. Child and Adolescent Health Measurement Initiative. Data Resource Center for Child and Adolescent Health website; 2007. [www.nschdata.org](http://www.nschdata.org).
5. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
6. Tax Extenders and Alternative Minimum Tax Relief Act of 2008, Pub. L. No. 110-343, Subtitle B (Paul Wellstone and Pete Domenici Mental Health Parity and Addiction Equity Act of 2008), [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=110\\_cong\\_public\\_laws&docid=f:publ343.110.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=110_cong_public_laws&docid=f:publ343.110.pdf).
7. Health Care and Education Reconciliation Act of 2010, Pub. L. No. 111-152, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ152.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ152.111.pdf).
8. Tolan PH, Dodge KA. Children's mental health as a primary care and concern: a system for comprehensive support and service. *Am Psychol*. Sep 2005;60(6):601-614.
9. Angold A, Costello EJ, Farmer EM, Burns BJ, Erkanli A. Impaired but undiagnosed. *J Am Acad Child Adolesc Psychiatry*. Feb 1999;38(2):129-137.
10. Briggs-Gowan MJ, Owens PL, Schwab-Stone ME, Leventhal JM, Leaf PJ, Horwitz SM. Persistence of psychiatric disorders in pediatric settings. *J Am Acad Child Adolesc Psychiatry*. Nov 2003;42(11):1360-1369.
11. Costello EJ, Angold A, Keeler GP. Adolescent outcomes of childhood disorders: the consequences of severity and impairment. *J Am Acad Child Adolesc Psychiatry*. Feb 1999;38(2):121-128.
12. Durlak JA, Wells AM. Primary prevention mental health programs for children and adolescents: a meta-analytic review. *Am J Community Psychol*. Apr 1997;25(2):115-152.
13. Durlak JA, Wells AM. Evaluation of indicated preventive intervention (secondary prevention) mental health programs for children and adolescents. *Am J Community Psychol*. Oct 1998;26(5):775-802.
14. National Scientific Council on the Developing Child. *Excessive stress disrupts the architecture of the developing brain: Working paper 3*;2005.
15. National Scientific Council on the Developing Child. *Mental health problems in early childhood can impair learning and behavior for life: Working paper 6*;2008.
16. Costello E, Shugart M. Above and below the threshold: Severity of psychiatric symptoms and functional impairment in a pediatric sample. *Pediatrics*. 1992;90(3):359-368.
17. Weisz JR, Sandler IN, Durlak JA, Anton BS. Promoting and protecting youth mental health through evidence-based prevention and treatment. *Am Psychol*. Sep 2005;60(6):628-648.
18. Halfon N, DuPlessis H, Inkelas M. Transforming the U.S. child health system. *Health Aff (Millwood)*. Mar-Apr 2007;26(2):315-330.
19. Rushton JL, Felt BT, Roberts MW. Coding of pediatric behavioral and mental disorders. *Pediatrics*. 2002;110:e8.
20. Hinshaw SP. The stigmatization of mental illness in children and parents: developmental issues, family concerns, and research needs. *J Child Psychol Psychiatry*. Jul 2005;46(7):714-734.
21. Crystal S, Olfson M, Huang C, Pincus H, Gerhard T. Broadened use of atypical antipsychotics: safety, effectiveness, and policy challenges. *Health Aff (Millwood)*. Sep-Oct 2009;28(5):w770-781.
22. Weissberg RP, Kumpfer KL, Seligman ME. Prevention that works for children and youth. An introduction. *Am Psychol*. Jun-Jul 2003;58(6-7):425-432.
23. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec.1301, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
24. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec.1201, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
25. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec.1302, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
26. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec.1311(j), [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
27. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec.1563(c)(4), [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
28. Cooper JL, Aratani Y, Knitzer J, et al. *Unclaimed children revisited: The status of children's mental health policy in the United States*. National Center for Children in Poverty;2008.
29. Brindis CD, Morreale MC, English A. The unique health care needs of adolescents. *The Future of Children*. 2003;13(1):117-135.
30. Aos S, Lieb R, Mayfield J, Miller M, Pennucci A. *Benefits and costs of prevention and early intervention programs for youth*. Olympia: Washington State Institute for Public Policy;2004.
31. Finch R, Sherrets D, Krackowsky K, Reagin A. *An employer's guide to child and adolescent mental health*: National Business Group on Health;2009.
32. McClellan JM, Werry JS. Evidence-based treatments in child and adolescent psychiatry: an inventory. *J Am Acad Child Adolesc Psychiatry*. Dec 2003;42(12):1388-1400.
33. Gottfredson D, Gottfredson G. Quality of school-based prevention programs: results from a national survey. *J Res Crime Delinq*. 2002;39(3).
34. Kazak AE, Hoagwood K, Weisz JR, et al. A meta-systems approach to evidence-based practice for children and adolescents. *Am Psychol*. Feb-Mar 2010;65(2):85-97.
35. Weiss B, Catron T, Harris V. A 2-year follow-up of the effectiveness of traditional child psychotherapy. *J Consult Clin Psychol*. Dec 2000;68(6):1094-1101.



36. Weiss B, Catron T, Harris V, Phung TM. The effectiveness of traditional child psychotherapy. *J Consult Clin Psychol*. Feb 1999;67(1):82-94.
37. Hoagwood K, Burns BJ, Kiser L, Ringeisen H, Schoenwald SK. Evidence-based practice in child and adolescent mental health services. *Psychiatr Serv*. Sep 2001;52(9):1179-1189.
38. Tynan WD, Stehl ML, Pendley JS. Health Insurance and Pediatric Psychology. In: Roberts MC, Steele RG, eds. *Handbook of Pediatric Psychology*. Fourth ed. NY: The Guilford Press; 2009:71-88.
39. Peele PB, Lave JR, Kelleher KJ. Exclusions and limitations in children's behavioral health care coverage. *Psychiatr Serv*. May 2002;53(5):591-594.
40. Flynn LM. Family perspectives on evidence-based practice. *Child Adolesc Psychiatr Clin N Am*. Apr 2005;14(2):217-224, vii.
41. Howell EM. Access to Children's Mental Health Services Under Medicaid and SCHIP. Urban Institute; August 31 2004.
42. Stroul B, Pires S, Armstrong M, et al. *RTC Study 3: Financing structures and strategies to support effective systems of care*. The Research and Training Center for Children's Mental Health;2009.
43. Raghavan R, Bright CL, Shadoin AL. Toward a policy ecology of implementation of evidence-based practices in public mental health settings. *Implement Sci*. 2008;3:26.
44. Bruns EJ, Hoagwood KE. State implementation of evidence-based practice for youths, Part I: responses to the state of the evidence. *J Am Acad Child Adolesc Psychiatry*. Apr 2008;47(4):369-373.
45. Bruns EJ, Hoagwood KE, Rivard JC, Wotring J, Marsenich L, Carter B. State implementation of evidence-based practice for youths, part II: recommendations for research and policy. *J Am Acad Child Adolesc Psychiatry*. May 2008;47(5):499-504.
46. Guevara JP, Feudtner C, Romer D, et al. Fragmented care for inner-city minority children with attention-deficit/hyperactivity disorder. *Pediatrics*. Oct 2005;116(4):e512-517.
47. Kim WJ. Child and adolescent psychiatry workforce: a critical shortage and national challenge. *Acad Psychiatry*. Winter 2003;27(4):277-282.
48. Kelleher KJ, Stevens J. Evolution of child mental health services in primary care. *Acad Pediatr*. Jan-Feb 2009;9(1):7-14.
49. Thomas CR, Holzer CE, 3rd. National distribution of child and adolescent psychiatrists. *J Am Acad Child Adolesc Psychiatry*. Jan 1999;38(1):9-15; discussion 15-16.
50. McKay MM, McCadam K, Gonzales JJ. Addressing the barriers to mental health services for inner city children and their caretakers. *Community Ment Health J*. Aug 1996;32(4):353-361.
51. Harrison ME, McKay MM, Bannon WM, Jr. Inner-city child mental health service use: the real question is why youth and families do not use services. *Community Ment Health J*. Apr 2004;40(2):119-131.
52. Blanchard LT, Gurka MJ, Blackman JA. Emotional, developmental, and behavioral health of American children and their families: A report from the 2003 National Survey of Children's Health. *Pediatrics*. 2006;117(6):e1202-1212.
53. Flisher AJ, Kramer RA, Grosser RC, et al. Correlates of unmet need for mental health services by children and adolescents. *Psychol Med*. Sep 1997;27(5):1145-1154.
54. Burns BJ, Phillips SD, Wagner HR, et al. Mental health need and access to mental health services by youths involved with child welfare: a national survey. *J Am Acad Child Adolesc Psychiatry*. Aug 2004;43(8):960-970.
55. McKay MM, Bannon WM, Jr. Engaging families in child mental health services. *Child Adolesc Psychiatr Clin N Am*. Oct 2004;13(4):905-921, vii.
56. Pescosolido BA, Jensen PS, Martin JK, Perry BL, Olafsdottir S, Fettes D. Public knowledge and assessment of child mental health problems: findings from the National Stigma Study-Children. *J Am Acad Child Adolesc Psychiatry*. Mar 2008;47(3):339-349.
57. Kazdin AE, Holland L, Crowley M. Family experience of barriers to treatment and premature termination from child therapy. *J Consult Clin Psychol*. Jun 1997;65(3):453-463.
58. Gilbody S, Bower P, Fletcher J, Richards D, Sutton A. Collaborative care for depression: A cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med*. 2006;166:2314-2321.
59. Kelleher KJ, Campo JV, Gardner WP. Management of pediatric mental disorders in primary care: where are we now and where are we going? *Curr Opin Pediatr*. Dec 2006;18(6):649-653.
60. Rushton J, Bruckman D, Kelleher K. Primary care referral of children with psychosocial problems. *Arch Pediatr Adolesc Med*. Jun 2002;156(6):592-598.
61. Williams J, Shore SE, Foy JM. Co-location of mental health professionals in primary care settings: three North Carolina models. *Clin Pediatr (Phila)*. Jul 2006;45(6):537-543.
62. Holt W. *The Massachusetts Child Psychiatry Access Project: Supporting mental health treatment in primary care*. The Commonwealth Fund;2010.
63. Guevara JP, Greenbaum PE, Shera D, Bauer L, Schwarz DF. Survey of mental health consultation and referral among primary care pediatricians. *Acad Pediatr*. Mar-Apr 2009;9(2):123-127.
64. Cummings NA, O'Donohue WT, Cummings JL. The financial dimension of integrated behavioral/primary care. *J Clin Psychol Med Settings*. Mar 2009;16(1):31-39.
65. Kautz C, Mauch D, Smith S. *Reimbursement of mental health services in primary care settings*. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration;2008. HHS Pub. no. SMA-08-4324.
66. AAP Task Force on Mental Health, AACAP Committee on Health Care Access and Economics. *White Paper: Improving mental health services in primary care: Reducing administrative and financial barriers to access and collaboration*;2009.
67. Patient Protection and Affordable Care Act of 2010, Pub. L. No. 111-148, Sec. 3021, 10306, [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111\\_cong\\_public\\_laws&docid=f:publ148.111.pdf](http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_public_laws&docid=f:publ148.111.pdf).
68. Bazelon Center for Mental Health Law. *Mix and Match: Using Federal Programs to Support Interagency Systems of Care for Children with Mental Health Care Needs*;2003.

## THE AUTHORS

**JANE E. KAVANAGH**, is senior research strategist with PolicyLab at The Children’s Hospital of Philadelphia Research Institute.

**ELIZABETH BROOKS, M.P.H., M.S.S.P.**, is a research associate with PolicyLab at The Children’s Hospital of Philadelphia Research Institute.

**SUSAN DOUGHERTY, Ph.D.**, is a psychologist and research scientist with PolicyLab at The Children’s Hospital of Philadelphia Research Institute.

**MARSHA GERDES, Ph.D.**, is senior psychologist with PolicyLab at The Children’s Hospital of Philadelphia Research Institute and clinical associate professor of pediatrics at the University of Pennsylvania.

**JAMES GUEVARA, M.D., M.P.H.**, is director of interdisciplinary initiatives with PolicyLab at The Children’s Hospital of Philadelphia Research Institute, an associate professor of pediatrics and epidemiology at the University of Pennsylvania, and an attending physician at The Children’s Hospital of Philadelphia.

**DAVID RUBIN, M.D., M.S.C.E.**, is director of PolicyLab at The Children’s Hospital of Philadelphia Research Institute, an associate professor of pediatrics at the University of Pennsylvania, and an attending physician at The Children’s Hospital of Philadelphia.

PolicyLab scientific co-director Cynthia Johnson Mollen, M.D., M.S.C.E., reviewed and approved this brief. PolicyLab’s Emily DePaul, Kathleen Noonan, Amanda O’Reilly, and Sarah Zlotnik provided editorial and content support.

*PolicyLab would like to thank early readers of this brief for their comments and advice.*

*Research for this project is supported with funds from The Children’s Hospital of Philadelphia and the Pew Charitable Trusts.*



The mission of PolicyLab at The Children’s Hospital of Philadelphia is to achieve optimal child health and well-being by informing program and policy changes through interdisciplinary research.

PolicyLab develops evidence-based solutions for the most challenging health-related issues affecting children. We partner with numerous stakeholders in traditional healthcare and other community locations to identify the programs, practices, and policies that support the best outcomes for children and their families. PolicyLab disseminates its findings beyond research and academic communities as part of its commitment to transform “evidence to action.”



---

*PolicyLab Evidence to Action briefs highlight PolicyLab research areas in the context of local and national policy issues to advance child health and well-being.*

[www.research.chop.edu/policylab](http://www.research.chop.edu/policylab)

### **PolicyLab**

The Children's Hospital of Philadelphia  
34th Street and Civic Center Boulevard  
CHOP North, Room 1528  
Philadelphia, PA 19104  
Phone: 267-426-5300  
Fax: 267-426-0380  
PolicyLab@email.chop.edu