

Incidence of Healthcare-associated Influenza-like Illness After a Primary Care Encounter Among Young Children

Date:

Mar 2018
Visit Article

Despite potential respiratory virus transmission in pediatric clinics, little is known about the risk of healthcareassociated viral infections attributable to outpatient encounters. We evaluated whether exposure to a pediatric clinic visit was associated with subsequent influenza-like illness (ILI). Using electronic health record data, we conducted a retrospective cohort study of all children aged <6 years who presented to a provider in a 29-clinic pediatric primary care network for a non-ILI-related encounter over 2 respiratory virus seasons (September 1, 2012, to April 30, 2014). We defined a risk period for potential healthcare-associated (HA) ILI of 1 to 8 days after a non-ILI clinic visit and identified all cases of ILI to compare the incidences of ILI visits 1 to 8 days after a non-ILI encounter and those of visits >8 days after a non-ILI encounter. Among 149987 children <6 years of age (mean age, 2.5 years) with ≥1 non-ILI visit during the study period, 531928 total encounters and 13951 (2.9%) ILI encounters were identified; 1941 (13.9%) occurred within the HA-ILI risk window. The incidence rate ratios (IRRs) for ILI 1 to 8 days after compared with ILI >8 days after a non-ILI visit during season 1 were 1.36 (95% confidence interval, 1.22-1.52) among children ≥2 years of age and 1.01 (95% confidence interval, 0.93-1.09) among children <2 years of age. Estimates remained consistent during season 2 and with a risk window of 3, 4, or 9 days. Pediatric clinic visits during a respiratory virus season were significantly associated with an increased incidence of subsequent ILI among children aged 2 to 6 years but not among those aged <2 years. These findings support the hypothesis that respiratory virus transmission in a pediatric clinic can result in HA ILI in young children.

Journal:

<u>Journal of the Pediatric Infectious Diseases Society</u>
Authors:

Feemster K, Localio R, Grundmeier R, Metlay JP, Coffin SE

Related Content

<u>Addressing Vaccine Hesitancy to Protect Children & Communities Against Preventable Diseases</u>

<u>Addressing Vaccine Hesitancy to Protect Children and Communities Against Preventable Diseases</u>