

---

## E-Cigarettes and Similar Devices

### Date:

Jan 2019

[Visit Article](#)

Electronic cigarettes (e-cigarettes) are the most commonly used tobacco product among youth. The 2016 US Surgeon General's Report on e-cigarette use among youth and young adults concluded that e-cigarettes are unsafe for children and adolescents. Furthermore, strong and consistent evidence finds that children and adolescents who use e-cigarettes are significantly more likely to go on to use traditional cigarettes—a product that kills half its long-term users. E-cigarette manufacturers target children with enticing candy and fruit flavors and use marketing strategies that have been previously successful with traditional cigarettes to attract youth to these products. Numerous toxicants and carcinogens have been found in e-cigarette solutions. Nonusers are involuntarily exposed to the emissions of these devices with secondhand and thirdhand aerosol. To prevent children, adolescents, and young adults from transitioning from e-cigarettes to traditional cigarettes and minimize the potential public health harm from e-cigarette use, there is a critical need for e-cigarette regulation, legislative action, and counterpromotion to protect youth.

### Journal:

[Pediatrics](#)

Authors:

Jenssen BP, Walley SC

### Related Content

[E-cigarettes, JUUL and Vaping: What Pediatric Health Care Providers Need to Know](#)

[E-Cigarettes: The New Face of Tobacco](#)

[Adolescents and E-cigarettes: Trends and What Pediatricians Need to Know](#)

[Tobacco Control and Treatment for the Pediatric Clinician: Practice, Policy, and Research Updates](#)