

Addressing Sexually Transmitted Infections in Pediatrics

Date:

Apr 2019

[Download Material](#)

Adolescents and young adults face a number of sexual and reproductive health needs that often go unmet. Youth ages 15–24 account for more than half of newly acquired sexually transmitted infections (STIs) each year in the United States. Many youth are unaware that they have a STI or other sexual or reproductive health condition; for example, only about half of HIV-infected adolescents are aware of their status. Another unmet adolescent sexual and reproductive health need relates to pregnancy; nearly 230,000 babies were born to teen mothers ages 15–19 in 2017 with three-quarters of teen pregnancies in 2011 being unintended.

Even though it is widely known that many youth struggle with sexual health challenges, pediatric practices don't always adequately support the sexual health needs of their patients due to a number of factors. In this *Research at a Glance* brief, Drs. Cynthia Mollen and Nadia Dowshen explain two different PolicyLab and Children's Hospital of Philadelphia studies that provide insight into the reality of screening for sexual health conditions in pediatric practices and one way we may be able to address some of the challenges to screening.

Authors:

Mollen C, Dowshen N

Topics

[Sexual & Reproductive Health](#)

Related Content

[Missed Opportunities for HIV Screening Among a Cohort of Adolescents With Recently Diagnosed HIV Infection in a Large Pediatric Hospital Care Network](#)

[Development of a Novel Computerized Clinical Decision Support System to Improve Adolescent Sexual Health Care Provision](#)

[Critical Access to Care: Bringing Contraception to Adolescents in Nontraditional Settings](#)

[How Expedited Partner Therapy Could Reduce Adolescent STD Rates in 200 Words](#)