

Social Support Networks Among Young Men and Transgender Women of Color Receiving HIV Pre-exposure Prophylaxis

Date:

Mar 2020

[Visit Article](#)

The aim of the study was to characterize perceived social support for young men and transgender women who have sex with men (YM/TWSM) taking HIV pre-exposure prophylaxis (PrEP). **METHODS:** Mixed-methods study of HIV-negative YM/TWSM of color prescribed oral PrEP. Participants completed egocentric network inventories characterizing their social support networks and identifying PrEP adherence support figures. A subset ($n = 31$) completed semistructured interviews exploring adherence support and qualities of PrEP support figures. We calculated proportions of role types (e.g., family), individuals disclosed to regarding PrEP use, and PrEP-supportive individuals within each participant network. Interviews were analyzed using an inductive approach. **RESULTS:** Participants ($n = 50$) were predominately African American men who have sex with men. Median age was 22 years (interquartile range: 20-23). Biologic family were the most common support figures, reported by 75% of participants (mean family proportion .37 [standard deviation (SD): .31]), followed by 67% reporting friends (mean friend proportion .38 [SD: .36]). Most network members were aware (mean disclosed proportion .74 [SD: .31]) and supportive (mean supportive proportion .87 [SD: .28]) of the participants' PrEP use. Nearly all (98%) participants identified ≥ 1 figure who provided adherence support; more often friends (48%) than family (36%). Participants characterized support as instrumental (e.g., transportation); emotional (e.g., affection); and social interaction (e.g., taking medication together). Key characteristics of PrEP support figures included closeness, dependability, and homophily (alikehood) with respect to sexual orientation. **CONCLUSIONS:** Although most YM/TWSM identified family in their support networks, friends were most often cited as PrEP adherence support figures. Interventions to increase PrEP adherence should consider integrated social network and family-based approaches.

Journal:

[Journal of Adolescent Health](#)

Authors:

Wood S, Dowshen N, Bauermeister JA, Lalley-Chareczko L, Franklin J, Petsis D, Swyrn M, Barnett K, Weissman GE, Koenig HC, Gross R

Related Content

[Comparing Different Measures of Retention in Care Among a Cohort of Adolescents and Young Adults Living with Behaviorally-acquired HIV](#)

[Barriers and Facilitators for Mental Healthcare in Pediatric Lupus and Mixed Connective Tissue Disease: A Qualitative Study of Youth and Parent Perspectives](#)

[Young Transgender Women's Attitudes Toward HIV Pre-exposure Prophylaxis](#)

[Sexual and Reproductive Health Considerations Among Transgender and Gender-expansive Youth](#)

[Family Characteristics in Sex Communication and Social Support: Implications for Emerging Adult Men Who Have Sex with Men's PrEP Engagement](#)

[Intersectionality and Health Inequities for Gender Minority Blacks in the U.S.](#)