

Association of Suicide Prevention Interventions With Subsequent Suicide Attempts, Linkage to Follow-up Care and Depression Symptoms for Acute Care Settings

[Visit Article](#)

IMPORTANCE: To prevent suicide deaths, acute care settings need tools to ensure individuals at risk of suicide access mental health care and remain safe until they do so. **OBJECTIVE:** To examine the association of brief acute care suicide prevention interventions with patients' subsequent suicide attempts, linkage to follow-up care, and depression symptoms at follow-up. **DATA SOURCES:** Ovid MEDLINE, Scopus, CINAHL, PsychINFO, Embase, and references of included studies using concepts of suicide, prevention, and clinical trial to identify relevant articles published January 2000 to May 2019. **STUDY SELECTION:** Studies describing clinical trials of single-encounter suicide prevention interventions were included. Two reviewers independently reviewed all articles to determine eligibility for study inclusion. **DATA EXTRACTION AND SYNTHESIS:** Two reviewers independently abstracted data according to PRISMA guidelines and assessed studies' risk of bias using the Cochrane Risk of Bias tool. Data were pooled for each outcome using random-effects models. Small study effects including publication bias were assessed using Peter and Egger regression tests. **MAIN OUTCOMES AND MEASURES:** Three primary outcomes were examined: subsequent suicide attempts, linkage to follow-up care, and depression symptoms at follow-up. Suicide attempts and linkage to follow-up care were measured using validated patient self-report measures and medical record review; odds ratios and Hedges *g* standardized mean differences were pooled to estimate effect sizes. Depression symptoms were measured 2 to 3 months after the encounter using validated self-report measures, and pooled Hedges *g* standardized mean differences were used to estimate effect sizes. **RESULTS:** A total of 14 studies, representing outcomes for 4270 patients, were included. Pooled-effect estimates showed that brief suicide prevention interventions were associated with reduced subsequent suicide attempts (pooled odds ratio, 0.69; 95% CI, 0.53-0.89), increased linkage to follow-up (pooled odds ratio, 3.04; 95% CI, 1.79-5.17) but were not associated with reduced depression symptoms (Hedges *g* = 0.28 [95% CI, ?0.02 to 0.59]). **CONCLUSIONS AND RELEVANCE:** In this meta-analysis, brief suicide prevention interventions were associated with reduced subsequent suicide attempts. Suicide prevention interventions delivered in a single in-person encounter may be effective at reducing subsequent suicide attempts and ensuring that patients engage in follow-up mental health care.

Journal:

[JAMA Psychiatry](#)

Authors:

Doupnik SK, Rudd B, Schmutte T, Worsley D, Bowden CF, McCarthy E, Eggan E, Bridge JA, Marcus SC

Topics

[Evidence-based Prevention, Screening & Treatment
Integration into Medical Settings](#)

Related Content

[Examining Hospitals' Role in Preventing Youth Suicide](#)

[Delivering Mental Health Care in Emergency and Hospital Settings in 200 Words](#)

[ED Visits and Readmissions After Follow-up for Mental Health Hospitalization](#)

[Adolescents' Experiences During "Boarding" Hospitalization While Awaiting Inpatient Psychiatric Treatment](#)

[Following Suicidal Ideation or Suicide Attempt](#)

[Trends in Pediatric Emergency Department Visits for Mental Health Conditions and Disposition by Presence of a Psychiatric Unit](#)

[Integrating Behavioral Health Services Into Medical Hospital Care for Children](#)

[Catching up to the Crisis: Opportunities for Pediatric Hospitals to Improve Children's Access to Mental Health Services](#)