

More Data, More Questions: No Simple Answer About Which Children Should Undergo Screening Neuroimaging for Clinically Occult Abusive Head Trauma

[Visit Article](#)

Abusive head trauma (AHT) is the leading cause of fatal child physical abuse. Victims may initially present with clinically occult AHT without overt signs of head trauma or with only subtle, nonspecific symptoms, which can make timely recognition of AHT challenging. Research has shown missed opportunities for early detection of AHT in the medical setting are common and can lead to repeated injury. Neuroimaging is needed to diagnose clinically occult AHT but is not without risk. Researchers have worked to understand the yield of neuroimaging in detection of clinically occult AHT and to identify risk factors, yet findings have varied widely across studies. Identifying which children undergoing physical abuse evaluations are at highest risk of clinically occult AHT is key to development of evidence-based imaging decision support tools for clinicians. Here we discuss the recent literature, identify potential reasons for variation across studies, and offer opportunities for future research.

Journal:

[Child Abuse & Neglect](#)

Authors:

Henry KM, Lindberg DM, Wood JN

Related Content

[Occult Head Injuries in Infants Evaluated for Physical Abuse](#)

[Impact of Child Abuse Clinical Pathways on Skeletal Survey Performance in High-risk Infants](#)

[Quality Improvement Initiative to Improve Abuse Screening Among Infants With Extremity Fractures](#)

[Advanced Cervical Spine Imaging in Abusive Head Trauma: An Update on Recent Literature and Future Directions](#)