

## COVID-19 and Changes in Child Obesity

### Date:

Mar 2021

[Visit Article](#)

The coronavirus disease 2019 (COVID-19) pandemic has caused economic hardship, school closings, shutdowns, limited physical activities, and increased food insecurity for many families. In December 2020, the American Academy of Pediatrics released interim guidance on how pediatricians can help children and families address healthy lifestyles and obesity management during the pandemic. To best support pediatric health system obesity prevention and management efforts, a better understanding of emerging disparities is needed to identify at-risk populations and develop focused interventions. Through a large pediatric primary care network, we evaluated the shifting rates of obesity for patients attending visits and explored disparities based on age, race and ethnicity, insurance, and income.

### Journal:

[Pediatrics](#)

Authors:

Jenssen BP, Kelly M, Powell M, Bouchelle Z, Mayne SL, Fiks AG

### Related Content

[Doing the Math: Addressing Social Determinants in Pediatric Obesity](#)

[The Research is Clear: Nurture Heart Health Early](#)

[When the Scales are Skewed: The Struggle for Equality in Child Weight Outcomes](#)

[Routine Cholesterol Tests and Subsequent Change in BMI among Overweight and Obese Children](#)

[How Have Rates of Childhood Obesity Changed Amid COVID-19 in 200 Words](#)

[Neighborhood Greenspace and Changes in Pediatric Obesity During COVID-19](#)

[Trends and Persistent Disparities in Child Obesity During the COVID-19 Pandemic](#)