
Association between Preferred Language and Risk of Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Children in the United States

Date:

Sep 2021

[Visit Article](#)

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has had a disproportionate impact on Black, Hispanic, and other individuals of color, although data on the effect of a person's language on SARS-CoV-2 infection are limited. Considering the barriers suffered by immigrants and non-English-speaking families, we tested whether children with a preferred language other than English was associated with SARS-CoV-2 infection. Children from families with a preferred language other than English had a higher predicted probability of SARS-CoV-2 test positivity (adjusted odds ratio, 3.76; 95% CI, 2.07-6.67) during the first wave of the pandemic. This discrepancy continued into the second wave (adjusted odds ratio, 1.64; 95% CI, 1.10-2.41), although the difference compared with families who prefer to speak English decreased over time. These findings suggest that children from non-English-speaking families are at increased risk of SARS-CoV-2 infection, and efforts to reverse systemic inequities causing this increased risk are needed.

Journal:

[American Journal of Tropical Medicine and Hygiene](#)

Authors:

Otto WR, Grundmeier RW, Montoya-Williams D, Njoroge WFM, Wallis KE, Gerber JS, Yun K

Topics

[Equitable Access to Services](#)

Related Content

[Addressing the Unique Needs of Immigrant and Limited-English-Proficient Communities During the COVID-19 Pandemic](#)

[How Health Systems Can Help Address Language Barriers to Achieve Digital Health Equity](#)

[COVID-19 Disparities and Impacts on Emotional Health & Well-being of Diverse Children](#)

[Are Refugee Essential Workers at a Higher Risk of Contracting COVID-19?](#)

[Disparities in SARS-CoV-2 Positivity Among Pregnant Patients with Limited English Proficiency](#)