

---

# Effects of Personalized Depression Prevention on Anxiety through 18-month Follow-up: A Randomized Controlled Trial

## Date:

Sep 2022

[Visit Article](#)

Depression and anxiety frequently co-occur and share several risk factors. There is some evidence for transdiagnostic effects of prevention programs on depression and anxiety. In the Personalized Depression Prevention (PDP) study, youth ( $n = 98$ ,  $M_{age} = 13.94$  years,  $SD = 1.67$ ) were classified as high or low on cognitive and interpersonal risk factors and randomized to either a cognitive-behavioral or an interpersonal prevention program. Some participants received a match between risk and prevention, others received a mismatch. Our initial work found evidence for the benefits of personalization on depression outcomes. In this paper, we focus on secondary anxiety outcomes through 18-months post-intervention. We found evidence for the benefits of personalized prevention on anxiety symptoms during the 18-month follow-up period, but not during the intervention. From post-intervention to 18-month follow-up matched youth showed a decrease in anxiety symptoms whereas mismatched youth showed a significant increase in symptoms ( $d = 0.87$ ,  $p = .001$ ). The rates of anxiety disorders were equivalent across the groups ( $p = 1.00$ ). Given the comorbidity of depression and anxiety, interventions that have effects on both may be an efficient and cost-effective approach to reducing the burden associated with these conditions. A risk-informed personalization approach to prevention may be one way to enhance the transdiagnostic effects of depression prevention.

## Journal:

[Behaviour Research and Therapy](#)

Authors:

Jones JD, Hankin BL, Gallop R, Haraden D, Sbrilli MD, Garber J, Young JF

## Topics

[Evidence-based Prevention, Screening & Treatment](#)