

# Use of Gastroesophageal Reflux Medications in Premature Infants After NICU Discharge

## Date:

Nov 2016

[Visit Article](#)

**OBJECTIVES:** To describe the epidemiology and management of gastroesophageal reflux (GER) medications started in the first year of life for premature infants.

**METHODS:** Retrospective review of a cohort of infants  $\leq 35$  weeks' gestation presenting for care by 168 days of age to a 30-site network between 2005 and 2009 ( $n = 2217$ ) and followed to 3 years of age. Medication frequency, types, and duration of use were assessed. Logistic regression identified factors associated with treatment.

**RESULTS:** Thirty-seven percent (812) were prescribed GER medications with 77% begun after NICU discharge. Ninety percent (727) received histamine-2 receptor antagonists, 33% (269) proton pump inhibitors, 22% (182) prokinetics; 40% (325) received  $>1$  medication. Outpatient medication was initiated at  $95 \pm 69$  days of life for total of  $294 \pm 249$  days (interquartile ratio: 117–359). Feeding issues (adjusted odds ratio [aOR] 2.05, 95% confidence interval [CI]: 1.24–3.39) were associated with outpatient initiation. Forty-three percent (322) of infants started before 6 months were still on at 1 year of age associated with gestational age  $<32$  weeks (aOR 1.76, 95% CI: 1.16–2.67), chronic lung disease (aOR 2.59, 95% CI: 1.29–5.22), and reactive airways disease (aOR 1.67, 95% CI: 1.05–2.65).

**CONCLUSIONS:** Of the 37% of the cohort on GER medications, 77% were started after NICU discharge with prolonged use of medications. Feeding difficulties were associated with starting medication and markers of chronic lung disease with continuation of treatment. With uncertain evidence of efficacy, use of these medications in a high-risk population should be carefully evaluated.

## Journal:

[Pediatrics](#)

Authors:

D'Agostino JA, Passarella M, Martin AE, Lorch SA

## Related Content

[A Decade of Improvement in Neonatal Intensive Care](#)

[Challenges to Measuring the Quality of Low-Risk Newborns](#)

[Perspectives of Low Socioeconomic Status Mothers of Premature Infants](#)

[Determining the Optimal Neonatal Care for Preterm Infants in the Era of Personalized Medicine](#)